

IDACIO® (adalimumab-aacf) Enrollment and Prescription Form

Required fields are marked with an asterisk (*).

1. Fax to KabiCare at 1-833-671-1010

2. Upload completed form file to kabicare.trialcard.com

3. Fax to patient's preferred specialty pharmacy Questions? Call 1.833.KABICARE (1.833.522.4227)

	Patient information	tion - to be c	, , , , , , , , , , , , , , , , , , , ,	, ,	ny datirorized per		•			
	First Name*:	Last N	ame*:		D.0	.B.* /	/	Gender:	М	F
	Address*:		(City*:			ate*:	Zip Code*	:	
	Phone Number*:	Home	Cell Email*	*•			iterpretat eeded:	ion services	Υ	N
	Parent/Guardian Name (if applicable)	C	aregiver Nam	1e (if applicable)		• • • • • • • • • • • • • • • • • • • •	anguage:			
	Best time to call (Monday-Friday):	Anytime	Morning	Afternoon	Evening	Check for	permission t	o leave messages	with caregiv	ver
	When did you start on treatment?*:	Not yet sta	rted 0-3	months ago	4-6 months ago	7-12 mg	nths ago	12+ mont	hs ago	
		Required	Patient Aut	horization and	l Additional Cor	sents				
	I have read and agree to the Patient Authorization to Share Personal Health Information and IDACIO® enrollment. (Section 1). I have read and agree to receive text messages and calls as explained in the Telephone Consumer Protection Act (TCPA) Consent (see section 2). IDACIO® Co-Pay Program: I have read and agree to the Terms and Conditions for participation (see section 3). Patient or authorized representative signature: I have read and agree to the Bridge to Commercial Coverage (see section 4 if applicable) and/or PAP if applicable (see section 4). I have read and agree to the Fair Credit Reporting Act (FCRA) Authorization (see section 5).									
	If authorized representative signature, explain au		ehalf of patient:					•	•	
-	Patient's Insurance Infor	rmation - Pro	ovide copies o	of the front and b	ack of all medica	l and prescr	iption ins	urance cards.		
	Beneficiary/Cardholder Name:				Prescription Insu					
	Medical Insurance:			Rx Group #:			Rx ID #:			
	Medical Insurance ID #:	Grou	ıp #:		Rx Bin #:		Rx PC			
			·	CARE PROVID	Commercial	Medicare		Medicaid	Unins	sured
	Dunganih ada Nama (Finah Lash)	V FU	R HEALIH	CARE PROVIL						
3	Prescriber's Name (First, Last)*:	Comboot Nor			Address:		Chahai	7: C		
	Office Phone:	Contact Nar			City:		State:	Zip Code	:	
	Contact Email:		ffice Fax*:		Tax ID:			PTAN:		
	NPI #*:	Rheumato	ology G	astroenterology	Dermatolog	y Oth	iei			
			Clini	cal Informatio	n					
	Diagnosis Date:		Clini	cal Informatio						
	Diagnosis Date: / /		Clini	Concom	itant Medications:					
4	Prior Therapies:	ſ		Concom Drug All	itant Medications:	Primar	w ICD-10 c	ode:		
4	Prior Therapies: Check all Rheumatoid arthritis indications Ankylosing spondylitis that apply: Juvenile idiopathic arthritis	tis in	Clini Plaque psoriasis Psoriatic arthritis	Concom Drug All Crohn's dise S Crohn's dise 6 years of a	itant Medications: ergies:		y ICD-10 c	ode:		
4	Prior Therapies: Check all indications that apply: Injection Rheumatoid arthritis Ankylosing spondylitis Juvenile idiopathic arthriting patients 2 years of age arthriting.	tis in nd older	Plaque psoriasis Psoriatic arthritis	Concom Drug All Crohn's disc S Crohn's disc 6 years of a Ulcerative of	itant Medications: ergies: ease in adults ease in pediatric patie ige and older colitis in adults	ents	-			
4	Prior Therapies: Check all indications that apply: Ankylosing spondylitis Juvenile idiopathic arthritipatients 2 years of age artificial I request supplemental injection Training:	tis in nd older ction training, for t	Plaque psoriasis >soriatic arthritis his patient. Order	Concom Drug All Crohn's disc s Crohn's disc 6 years of a Ulcerative o	itant Medications: ergies: ease in adults ease in pediatric patie ige and older colitis in adults	ents pharmacy pres	cription belo			
4 5	Prior Therapies: Check all indications that apply: Ankylosing spondylitis Juvenile idiopathic arthritipatients 2 years of age artificial I request supplemental injection Training:	tis in nd older ction training, for t escription - Se	Plaque psoriasis >soriatic arthritis his patient. Order	Concom Drug All Crohn's disc s Crohn's disc 6 years of a Ulcerative o	itant Medications: ergies: ease in adults ease in pediatric pation ige and older colitis in adults ar. Fill out and sign the	ents pharmacy pres	cription belo			
4	Prior Therapies: Check all indications that apply: Injection Training: Pharmacy Prepatient's preferred specialty pharmacy: Starting Rheumatoid arthritis Ankylosing spondylitis Juvenile idiopathic arthritic patients 2 years of age and arthritic patients 2	tis in nd older ction training, for t escription - Se	Plaque psoriasis Psoriatic arthritis his patient. Order Plect medicati	Concom Drug All Crohn's disc S Crohn's disc 6 years of a Ulcerative of the control of the c	itant Medications: ergies: ease in adults ease in pediatric patie age and older colitis in adults ar. Fill out and sign the ign corresponding	pharmacy pres prescription	cription belo			
4	Prior Therapies: Check all indications that apply: Injection Training: Pharmacy Preserved Specialty pharmacy:	tis in and older ction training, for the cescription - Secription - Se	Plaque psoriasis Psoriatic arthritis his patient. Order Plect medicati	Concom Drug All Crohn's disc S Crohn's disc 6 years of a Ulcerative of a valid for up to one years ion, fill out and s	itant Medications: ergies: ease in adults ease in pediatric patie ge and older colitis in adults ar. Fill out and sign the ign corresponding	pharmacy pres g prescription OR PEN IDA old CIO Citrate-	on below.			
4	Prior Therapies: Check all indications that apply: Injection Training: Pharmacy Prepared Specialty pharmacy: Starting AUTOINJECTOR PETHORNOOSE 1 SYRINGE IDACIO City Presentation 40 mg/0.8 mL Adult Crohn's Disease, Ulcerative Colitis, and (40 kg (88lbs) and greater) SIG 160 mg SC inj. on Day 1 (given in one)	tis in and older ction training, for the cascription - Section training for the cascription - Section - Section training for the cascription - Section training for the cascription - Section	Plaque psoriasis Psoriatic arthritis his patient. Order elect medicati ate-Free phn's Disease two consecutive	Concom Drug All Crohn's disc S Crohn's disc 6 years of a Ulcerative of valid for up to one year on, fill out and s Ongoing therapy: Choose 1 Presentation SIG days);	itant Medications: ergies: ease in adults ease in pediatric patie ge and older colitis in adults ear. Fill out and sign the ign corresponding AUTOINJECT 40 mg/0.8 m SYRINGE IDA 40 mg/0.8 m 40 mg/0.8 m 40 mg/0.8 m	pharmacy pres g prescription OR PEN IDA IL CIO Citrate- IL ery other wee ery week†	on below. CIO Citrat			
4	Prior Therapies: Check all indications that apply: Injection Training: Pharmacy Preserved specialty pharmacy: Starting AUTOINJECTOR PE 40 mg/0.8 mL Choose 1 SYRINGE IDACIO Cipresentation 40 mg/0.8 mL Adult Crohn's Disease, Ulcerative Colitis, and (40 kg (88lbs) and greater) SIG 160 mg SC inj. on Day 1 (given in one 80 mg SC inj. on Day 15; and 40 mg #QS No Refills	tis in and older ction training, for the cascription - Section training for the cascription - Section training for the cascription training for	Plaque psoriasis Psoriatic arthritis his patient. Order elect medicati ate-Free ohn's Disease two consecutive her week startin	Concom Drug All Crohn's disc S Crohn's disc 6 years of a Ulcerative of valid for up to one year on, fill out and s Ongoing therapy: Choose 1 Presentation SIG days); ag on Day 29.	itant Medications: ergies: ease in adults ease in pediatric patie ge and older colitis in adults er. Fill out and sign the ign corresponding AUTOINJECT 40 mg/0.8 m SYRINGE IDA 40 mg/0.8 m	pharmacy pres prescription OR PEN IDA L CIO Citrate- L ery other wee ery week [†] ery other wee	cription below. CIO Citrat Free k			
4	Prior Therapies: Check all indications that apply: Injection Training: Pharmacy Preserved specialty pharmacy: Starting AUTOINJECTOR PE 40 mg/0.8 mL Choose 1 SYRINGE IDACIO Ci 40 mg/0.8 mL Adult Crohn's Disease, Ulcerative Colitis, a (40 kg (88lbs) and greater) SIG 160 mg SC inj. on Day 1 (given in one 80 mg SC inj. on Day 15; and 40 mg	tis in and older ction training, for the ction training training training spoint training tr	Plaque psoriasis Psoriatic arthritis his patient. Order elect medicati ate-Free ohn's Disease two consecutive her week startin	Concom Drug All Crohn's disc S Crohn's disc 6 years of a Ulcerative of valid for up to one year on, fill out and s Ongoing therapy: Choose 1 Presentation SIG days); ag on Day 29.	itant Medications: ergies: erase in adults ease in pediatric patie ige and older colitis in adults er. Fill out and sign the ign corresponding AUTOINJECT 40 mg/0.8 m SYRINGE IDA 40 mg/0.8 m	pharmacy pres prescription OR PEN IDA L CIO Citrate- L ery other wee ery week [†] ery other wee h 3 m	cription below. CIO Citrat Free k	w. - e-Free Refills:		
4	Prior Therapies: Check all indications that apply: Injection Training: Pharmacy Pre Patient's preferred specialty pharmacy: Starting AUTOINJECTOR PE 40 mg/0.8 mL Choose 1 SYRINGE IDACIO Cide Presentation 40 mg/0.8 mL Adult Crohn's Disease, Ulcerative Colitis, and (40 kg (88lbs) and greater) SIG 160 mg SC inj. on Day 1 (given in one 80 mg SC inj. on Day 15; and 40 mg #QS No Refills Rheumatoid Arthritis, Psoriatic Arthritis, Idiopathic Arthritis (30 kg (66 lbs) and greater) SIG 40 mg SC inj. week 1 #QS No Refills Plaque Psoriasis	tis in and older ction training, for the ction training training training spoint training tr	Plaque psoriasis Psoriatic arthritis his patient. Order elect medicati ate-Free ohn's Disease two consecutive her week startin	Concom Drug All Crohn's disc S Crohn's disc 6 years of a Ulcerative of valid for up to one year on, fill out and s Ongoing therapy: Choose 1 Presentation SIG days); ag on Day 29.	itant Medications: ergies: erase in adults ease in pediatric patie ige and older colitis in adults er. Fill out and sign the ign corresponding AUTOINJECT 40 mg/0.8 m SYRINGE IDA 40 mg/0.8 m	pharmacy pres prescription OR PEN IDA L CIO Citrate- L ery other wee ery week [†] ery other wee h 3 m	cription below. CIO Citrat Free k	w. - e-Free Refills:		
4	Prior Therapies: Check all indications that apply: Injection Training: Pharmacy Pre Patient's preferred specialty pharmacy: Starting AUTOINJECTOR PE 40 mg/0.8 mL Choose 1 SYRINGE IDACIO Cide Presentation 40 mg/0.8 mL Adult Crohn's Disease, Ulcerative Colitis, and (40 kg (88lbs) and greater) SIG 160 mg SC inj. on Day 1 (given in one 80 mg SC inj. on Day 15; and 40 mg #QS No Refills Rheumatoid Arthritis, Psoriatic Arthritis, Idiopathic Arthritis (30 kg (66 lbs) and greater) SIG 40 mg SC inj. week 1 #QS No Refills	tis in and older ction training, for the ction training training training spoint training tr	Plaque psoriasis Psoriatic arthritis his patient. Order elect medicati ate-Free ohn's Disease two consecutive her week startin	Concom Drug All Crohn's disc S Crohn's disc 6 years of a Ulcerative of valid for up to one year on, fill out and s Ongoing therapy: Choose 1 Presentation SIG days); ag on Day 29.	itant Medications: ergies: erase in adults ease in pediatric patie ige and older colitis in adults er. Fill out and sign the ign corresponding AUTOINJECT 40 mg/0.8 m SYRINGE IDA 40 mg/0.8 m	pharmacy pres prescription OR PEN IDA L CIO Citrate- L ery other wee ery week [†] ery other wee h 3 m	cription below. CIO Citrat Free k	w. - e-Free Refills:		
4	Prior Therapies: Check all indications that apply: Injection Training: Pharmacy Pre Patient's preferred specialty pharmacy: Starting AUTOINJECTOR PE 40 mg/0.8 mL Choose 1 SYRINGE IDACIO Ci Presentation 40 mg/0.8 mL Adult Crohn's Disease, Ulcerative Colitis, a (40 kg (88lbs) and greater) SIG 160 mg SC inj. on Day 1 (given in one 80 mg SC inj. on Day 15; and 40 mg #QS No Refills Rheumatoid Arthritis, Psoriatic Arthritis, Idiopathic Arthritis (30 kg (66 lbs) and greater) SIG 40 mg SC inj. week 1 #QS No Refills Plaque Psoriasis SIG 80 mg SC inj. week 1	tis in ad older ction training, for the escription - Secription - Sec	Plaque psoriasis Psoriatic arthritis his patient. Order elect medicati ate-Free two consecutive her week startin adylitis, and Jun is medically nece	Concom Drug All Crohn's disc S Crohn's disc 6 years of a Ulcerative of valid for up to one year on, fill out and s Ongoing therapy: Choose 1 Presentation SIG days); ng on Day 29. venile	itant Medications: ergies: ease in adults ease in pediatric patie ige and older colitis in adults ear. Fill out and sign the ign corresponding AUTOINJECT 40 mg/0.8 m SYRINGE IDA 40 mg/0.8 m 40 mg/0.8 m 40 mg/0.8 m Omyon SC inj. ev 40 mg/0.8 m Tosage frequency is	pharmacy pres g prescription OR PEN IDA old CIO Citrate- old ery other wee ery week† ery other wee h 3 m recommended of	cription below. CIO Citrat Free k ch ponths ponths	e-Free Refills:	mTX.	





IDACIO® (adalimumab-aacf) Enrollment and Prescription Form

- 1. Fax to KabiCare at 1-833-671-1010
- 2. Upload completed form file to kabicare.trialcard.com
- 3. Fax to patient's preferred specialty pharmacy Questions? Call 1.833.KABICARE (1.833.522.4227)

Important Safety Information

SERIOUS INFECTIONS

Patients treated with IDACIO® (adalimumab-aacf) are at increased risk for developing serious infections that may lead to hospitalization or death. Most patients who developed these infections were taking concomitant immunosuppressants such as methotrexate or corticosteroids.

Discontinue IDACIO® if a patient develops a serious infection or sepsis.

Reported infections include:

- Active tuberculosis (TB), including reactivation of latent TB. Patients with TB have frequently presented with disseminated or extrapulmonary disease. Test patients for latent TB before IDACIO® use and during therapy. Initiate treatment for latent TB prior to IDACIO® use.
- Invasive fungal infections, including histoplasmosis, coccidioidomycosis, candidiasis, aspergillosis, blastomycosis, and pneumocystosis. Patients with histoplasmosis or other invasive fungal infections may present with disseminated, rather than localized, disease. Antigen and antibody testing for histoplasmosis may be negative in some patients with active infection. Consider empiric antifungal therapy in patients at risk for invasive fungal infections who develop severe systemic illness.
- Bacterial, viral and other infections due to opportunistic pathogens, including Legionella and Listeria

Carefully consider the risks and benefits of treatment with IDACIO® prior to initiating therapy in patients: 1. with chronic or recurrent infection, 2. who have been exposed to TB, 3. with a history of opportunistic infection, 4. who resided in or traveled in regions where mycoses are endemic, 5. with underlying conditions that may predispose them to infection. Monitor patients closely for the development of signs and symptoms of infection during and after treatment with IDACIO®, including the possible development of TB in patients who tested negative for latent TB infection prior to initiating therapy.

Treatment with IDACIO® should not be initiated in patients with an active infection, including localized infections.

Patients 65 years of age and older, patients with co-morbid conditions and/or patients taking concomitant immunosuppressants (such as corticosteroids or methotrexate), may be at greater risk of infection.

Discontinue IDACIO® if a patient develops a serious infection or sepsis. For a patient who develops a new infection during treatment with IDACIO®, closely monitor them, perform a prompt and complete diagnostic workup appropriate for an immunocompromised patient, and initiate appropriate antimicrobial therapy.

Drug interactions with biologic products: A higher rate of serious infections has been observed in RA patients treated with rituximab who received subsequent treatment with a TNF blocker. An increased risk of serious infections has been seen with the combination of TNF blockers with anakinra or abatacept, with no demonstrated added benefit in patients with RA. Concomitant administration of adalimumab products with other biologic DMARDs (e.g., anakinra or abatacept) or other TNF blockers is not recommended based on the possible increased risk for infections and other potential pharmacological interactions.

MALIGNANCY

Lymphoma and other malignancies, some fatal, have been reported in children and adolescent patients treated with TNF blockers, including adalimumab products. Post-marketing cases of hepatosplenic T-cell lymphoma (HSTCL), a rare type of T-cell lymphoma, have been reported in patients treated with TNF blockers including adalimumab products. These cases have had a very aggressive disease course and have been fatal. The majority of reported TNF blocker cases have occurred in patients with Crohn's disease or ulcerative colitis and the majority were in adolescent and young adult males. Almost all these patients had received treatment with azathioprine or 6-mercaptopurine concomitantly with a TNF blocker at or prior to diagnosis. It is uncertain whether the occurrence of HSTCL is related to use of a TNF blocker or a TNF blocker in combination with these other immunosuppressants

- Consider the risks and benefits of TNF-blocker treatment prior to initiating or continuing therapy in a patient with known malignancy.
- In clinical trials of some TNF-blockers, including adalimumab products, more cases of malignancies were observed among TNF-blocker-treated patients compared to control patients.
- Non-melanoma skin cancer (NMSC) was reported during clinical trials for adalimumabtreated patients. Examine all patients, particularly those with a history of prolonged immunosuppressant or PUVA therapy, for the presence of NMSC prior to and during treatment with IDACIO®.
- In adalimumab clinical trials, there was an approximate 3-fold higher rate of lymphoma than expected in the general U.S. population. Patients with chronic inflammatory diseases, particularly those with highly active disease and/or chronic exposure to immunosuppressant therapies, may be at higher risk of lymphoma than the general population, even in the absence of TNF blockers.
- Postmarketing cases of acute and chronic leukemia were reported with TNF blocker use.
 Approximately half of the postmarketing cases of malignancies in children, adolescents, and young adults receiving TNF blockers were lymphomas; other cases included rare malignancies associated with immunosuppression and malignancies not usually observed in children and adolescents.

HYPERSENSITIVITY

Anaphylaxis and angioneurotic edema have been reported following administration of adalimumab products. If an anaphylactic or other serious allergic reaction occurs, immediately discontinue administration of IDACIO® and institute appropriate therapy. In clinical trials of adalimumab, hypersensitivity reactions (e.g., rash, anaphylactoid reaction, fixed drug reaction, non-specified drug reaction, urticaria) have been observed.

HEPATITIS B VIRUS REACTIVATION

Use of TNF blockers, including IDACIO®, may increase the risk of reactivation of hepatitis B virus (HBV) in patients who are chronic carriers of this virus. In some instances, HBV reactivation occurring in conjunction with TNF blocker therapy has been fatal.

Evaluate patients at risk for HBV infection for prior evidence of HBV infection before initiating TNF blocker therapy.

Exercise caution in prescribing TNF blockers for patients identified as carriers of HBV. In patients who develop HBV reactivation, stop IDACIO® and initiate effective anti-viral therapy with appropriate supportive treatment. The safety of resuming TNF blocker therapy after HBV reactivation is controlled is not known. Therefore, exercise caution when considering resumption of IDACIO® therapy in this situation and monitor patients closely.

NEUROLOGICAL REACTIONS

Use of TNF blocking agents, including adalimumab products, has been associated with rare cases of new onset or exacerbation of clinical symptoms and/or radiographic evidence of central nervous system demyelinating disease, including multiple sclerosis (MS) and optic neuritis, and peripheral demyelinating disease, including Guillain-Barré syndrome.

Exercise caution in considering the use of IDACIO® in patients with pre-existing or recentonset central or peripheral nervous system demyelinating disorders; discontinuation of IDACIO® should be considered if any of these disorders develop.

HEMATOLOGICAL REACTIONS

Rare reports of pancytopenia, including aplastic anemia, have been reported with TNF blockers. Medically significant cytopenia has been infrequently reported with adalimumab products.

Consider stopping IDACIO® if significant hematologic abnormalities occur.

CONGESTIVE HEART FAILURE

Worsening or new onset congestive heart failure (CHF) may occur; exercise caution and monitor carefully.

AUTOIMMUNITY

Treatment with adalimumab products may result in the formation of autoantibodies and, rarely, in development of a lupus-like syndrome. Discontinue treatment if symptoms of a lupus-like syndrome develop.

IMMUNIZATIONS

Patients on IDACIO® should not receive live vaccines. Pediatric patients, if possible, should be brought up to date with all immunizations before initiating IDACIO® therapy. The safety of administering live or live-attenuated vaccines in infants exposed to adalimumab products in utero is unknown. Risks and benefits should be considered prior to vaccinating (live or live-attenuated) exposed infants.

ADVERSE REACTIONS

The most common adverse reactions in adalimumab clinical trials ($^{10\%}$) were: infections (e.g. upper respiratory, sinusitis), injection site reactions, headache and rash.

INDICATIONS

- Rheumatoid Arthritis (RA): reducing signs and symptoms, inducing major clinical response, inhibiting the progression of structural damage, and improving physical function in adult patients with moderately to severely active RA.
- Juvenile Idiopathic Arthritis (JIA): reducing signs and symptoms of moderately to severely active polyarticular JIA in patients 2 years of age and older.
- Psoriatic Arthritis (PsA): reducing signs and symptoms, inhibiting the progression of structural damage, and improving physical function in adult patients with active PsA.
- Ankylosing Spondylitis (AS): reducing signs and symptoms in adult patients with active AS.
- Crohn's Disease (CD): treatment of moderately to severely active Crohn's disease in adults and pediatric patients 6 years of age and older.
- Ulcerative Colitis (UC): treatment of moderately to severely active ulcerative colitis in adult patients.
 <u>Limitations of Use</u>: Effectiveness has not been established in patients who have lost response to or were intolerant to TNF blockers.
- Plaque Psoriasis (Ps): treatment of adult patients with moderate to severe chronic plaque psoriasis who are candidates for systemic therapy or phototherapy, and when other systemic therapies are medically less appropriate.

Click to see full Prescribing Information, including **Boxed WARNING,** for IDACIO® (adalimumab-aacf).





IDACIO® (adalimumab-aacf) Enrollment and Prescription Form

- 1. Fax to KabiCare at 1-833-671-1010
- 2. Upload completed form file to kabicare.trialcard.com
- 3. Fax to patient's preferred specialty pharmacy Questions? Call 1.833.KABICARE (1.833.522.4227)

ADDITIONAL TERMS AND CONDITIONS FOR ENROLLMENT

Section 1: Patient Authorization for the Use and Disclosure of Protected Health Information

PATIENT AUTHORIZATION AND RELEASE TO COLLECT, USE, AND DISCLOSE HEALTH INFORMATION

By my signature above, I agree to allow my doctors, pharmacies, specialty pharmacy(ies), and health insurers (collectively "Healthcare Providers"), to use and disclose my personal health information related to this enrollment form or my use or potential use of IDACIO to Fresenius Kabi and its agents, authorized representatives, and contractors, including TrialCard Incorporated, as further described below

Information to Be Disclosed: My protected health information (as such term is defined in the Health Insurance Portability and Accountability Act of 1996 ("HIPAA") and regulations thereunder), as well as other state and/or federally protected personal information, including my personal contact and other demographic information, all medical records and financial information, and information relating to my treatment, the coordination of my treatment, and the delivery, packaging, and recipt of certain medication prescribed to me (collectively, my "Information"). "Information").

Persons to Whom My Information May Be Disclosed: Fresenius Kabi and the KabiCare Patient Support Program, including any third parties responsible for the administration of the KabiCare Patient Support Program.

Purposes for Which the Disclosures Are to Be Made:
Disclosures of my Information may be made to KabiCare
Patient Support Program so that KabiCare Patient
Support Program may use and disclose my Information
for purposes of:
1) Communicating with my Healthcare Providers
about my prescription and medical condition, includin
to facilitate the order, fulfillment and delivery of my

- to facilitate the order, fulfillment and delivery of my prescription as needed; Establishing my eligibility for benefits from my health plan or other programs; Contacting my insurer on my behalf to determine if I am eligible for health insurance coverage or other funds; Providing appropriate product and reimbursement support:

- 4) Providing appropriate product and reimbursement support;
 5) Contacting me regarding case management and/ or educational information or training offered by or through the KabiCare Patient Support Program;
 6) Contacting me regarding this Authorization or my use or potential use of my prescriptions and providing me with related communications, including through messages left for me that disclose that I take or may take certain prescription medications;
 7) Contacting me to administer, evaluate, and improve the KabiCare Patient Support Programs, including analyzing the usage patterns and the effectiveness of services and helping to develop new products, services, and programs, and for other general business and administrative purposes; and
 8) Disclosing my Information to third parties if required by law.

- law.
 y signing this Authorization, I acknowledge my olderstanding that:
 I understand that I may refuse to sign this
 Authorization and that my refusal to sign this
 Authorization will not affect my right to treatment or payment of benefits for health care. I understand that if I refuse to sign, I will not be alicible to regions ungent through the
- care. I understand that if I refuse to sign, I will note eligible to receive support through the KabiCare Patient Support Program. I have the right to revoke this Authorization at any time by mailing 2250 Perimeter Park Dr., Suite 300 Morrisville, NC 27560. Revoking this Authorization will prohibit further uses and disclosures of my Information by the KabiCare Patient Support Program, except to the extent those uses and disclosures have been made in reliance on this Authorization and as permitted by applicable law.
- reliance on this Authorization and as permitted by applicable law. Certain pharmacy providers or other Healthcare Providers may receive remuneration for the use or disclosure of my Information, as permitted by this Authorization.
- or disclosure of my Information, as permitted by this Authorization.

 Once my Information is released to KabiCare Patient Support Program based on this Authorization, my Information may not be subject to all of the protections and safequards provided by HIPAA or other federal and state privacy laws. However, I understand that KabiCare Patient Support Program has agreed to use or disclose my Information received only for the purposes described in this Authorization or as required by law.

 This Authorization will remain in effect for a period of two (2) years after I sign it unless a shorter period is required by state law or is revoked by me earlier in writing, I am entitled to receive a copy of this signed Authorization.

Authorization. The patient, or the patient's authorized representative, Ine patient, or the patient's authorized representative, MUST sign this form to participate in the KabiCare Patient Support Program. My signature, signifying my agreement with this Authorization, is provided on page 1 of this Enrollment Form, where it states, 'I have read and agree to the Patient Authorization for the Use and Disclosure of Protected Health Information (section 1).' If an authorized representative signs for the patient, please indicate relationship to the patient.

Section 2: Telephone Consumer Protection Act (TCPA)

By checking the first box in the "Required Patient By Checking the invist box in the required radius Authorization and Additional Consents" section on page 1, you understand that the personal information you provide will be shared with KabiCare, and its third-party partners, including IrailCard Incorporated, You also attorize KabiCare, and its partners, to communicate with you about products, health conditions, copay and financial assistance. You are providing consent that you can be contacted by mail, email, phone or autodialed text messages under the guidelines of the Telephone Consumer Protection Act at the phone number you provide. You may opt out of individual communications from KabiCare or its partners entirely at any time by calling 1+833+KabiCare. KabiCare and companies providing services to KabiCare will not sell or rent your personally identifiable information, as described in the Privacy Policy. For clarity, phone contact includes telephonic contact for the purposes of advertising or telemarketing and may include either or both of (1) autodialed prerecorded voice calls, and (2) autodialed autodialed prerecorded voice calls, and (2) autodialed artificial voice calls.

You understand that you are not required to agree to receive any phone, mail, or text messages as a condition of participation in this Program, that message and data rates may apply depending on your phone carrier, and you may stop receiving text messages at any time by texting the word STOP in response to a text received from KabiCare or its nathrate.

Section 3: KabiCare Copay Assistance Program TERMS & CONDITIONS

To receive benefits under the Copay Assistance Program, the patient may contact the KabiCare Patient Support Program for current Program Product(s) subject to these Terms and Conditions. By participating in the Copay Assistance Program, patient acknowledges and agrees that he/she is eligible to participate and that he/she understands and agrees to comply with these Terms and Conditions.

- Conditions.

 Patient must be prescribed the Program Product for an FDA-approved indication.

 Patient must have commercial (private or non-governmental) health insurance that provides coverage for the cost of the Program Product under a pharmacy or medical benefit
- plan. The Copay Assistance Program is valid for patients who have a valid prescription for a Fresenius Kabi medication and who are not reimbursed for the entire cost of the prescription by their commercial insurance plan. The Copay Assistance Program is not valid for patients enrolled in Medicaid, Medicare (including a Medicare Part D or Medicare Advantage plan, a Medigap plan, or an employer-sponsored health plan or prescription drug benefit program for Medicare-eligible retirees), Veteran Affairs health care programs, Department of Defense health care programs, TRICARE, CHAMPUS, Puerto Rico Government Health Insurance Plan, or any other state or federal medical or Plan , or any other state or federal medical or pharmaceutical benefit program or pharmaceutical assistance program, including pharmaceutical assistance program, including any state pharmaceutical assistance programs (collectivelty, "Government Programs"). Patients who move from commercial insurance to Government Programs will no longer be eligible to participate in the Copay Assistance Program and agree to notify the Copay. If the patient lives in Massachusetts, the Copay Assistance Program expires on the earlier of: (i) the Expiration Date set Forth below, (ii) the date an AB-rated generic equivalent becomes available for the Program Product; or (iii) January 31, 2026, absent a change in Massachusetts state law. If the patient lives in California, the Copay Assistance Program expires on the earlier of: (ii)
- If the patient lives in California, the Copay Assistance Program expires on the earlier of: (i) the Expiration Date set forth below; or (ii) the date an FDA approved therapeutically equivalent for the Program Product or over the counter product with the same active ingredients becomes available. Patients must have an out-of-pocket cost for the Program Product prior to the Expiration Date of the Copay Assistance Program. The benefit available under the Copay Assistance Program. The the patient's private health insurance company indicates that the patient is obligated to pay for up to a per that the patient is obligated to pay for up to a per
- that the patient is obligated to pay for up to a per
- that the patient is obligated to pay for up to a per syringe/annual maximum. The Program may apply to patient out-of-pocket costs incurred for Program Product subject to per syringe/annual maximums based on Program Product administration date. After reaching the maximum Copay Assistance Program benefit, the patient will be responsible for all remaining out-of-pocket expenses. The patient or provider may contact the KabiCare Patient Support Program for more information. This Copay Assistance Program benefit is for commercially insured patients only. Uninsured and cash paying patients may be eligible for other types of support not part of the Copay Assistance Program.
- Assistance Program. The Patient and participating pharmacy
- The Patient and participating pharmacy or healthcare professional agree not to seek reimbursement for all, or any part, of the benefit received by the patient through the Copay Assistance Program. Participating patients and pharmacies or health care professionals are responsible for reporting receipt of Copay Assistance Program benefits as may be required by law. Patient must be a resident of the United States or the Commonwealth of Puerto Rico. Product must originate and be administered to patient in the United States or the Commonwealth of Puerto Rico. All information applicable to the Copay Assistance Program requested on the Kabilcare.US site must
- All miorimation applicable to the Copay Assistance Program requested on the KablCare.US site must be provided, and all certifications must be provided. No other purchase is necessary.
 The Copay Assistance Program is not insurance. It is illegal to sell, purchase, trade, counterfeit, or

- duplicate, or offer to sell, purchase, trade counterfeit, or duplicate the Copay Assistance Program card. Void
- if reproduced. The Copay Assistance Program is intended to comply

- The Copay Assistance Program is intended to comply with all applicable laws and regulations, including, without limitation, the federal Anti-Kickback Statute, its implementing regulations, and related guidance interpreting the federal Anti-Kickback Statute. The Copay Assistance Program is void where prohibited by law, taxed, or restricted. The Copay Assistance Program is not transferable. No substitutions are permitted. The Copay Assistance Program benefit has no cash value and cannot be combined with any other Copay Assistance Program, free trial, discount, rebate, prescription savings card, or other offer. The full value of the Copay Assistance Program benefit is intended to pass entirely to the patient to other individual or entity is entitled to receive any discount or other amount in connection with the Copay Assistance Program.
- Copay Assistance Program. This offer is not conditioned on any past, present, or future purchase obligation, and the Copay Assistance Program does not obligate the use of any specific

- Program does not obligate the use of any specific product or provider. To the extent applicable, this offer will be accepted only at participating pharmacies.

 KabiCare reserves the right to rescind, revoke, terminate, or amend the Copay Assistance Program at any time without notice.

 Data related to patient's receipt of Copay Assistance Program benefits may be collected, analyzed, and shared with KabiCare, for market research and other purposes related to assessing Copay Assistance Programs Data shared with KabiCare, for market research and other purposes related to assessing Copay Assistance Programs. Data shared with KabiCare will be aggregated and de-identified, meaning it will be combined with data related to other Copay Assistance Program redemptions and will not identify patient. The Terms and Conditions of the Copay Assistance Program are valid for Program Product only, and Fresenius Kabi reserves the right to rescind, revoke, or amend the Program without notice.

Section 4: Bridge to Commercial Coverage - Patient

KabiCare provides a limited and temporary supply of free product through the KabiCare Commercial Bridge Progra ("Commercial Bridge Program") for eligible commercially insured patients when a prior authorization request has been pending with the payer for more than 7 days and when other program eligibility criteria have been satisfied.

The patient and participating providers may not seek reimbursement for any free product provided under the Commercial Bridge Program nor does the Commercial Bridge Program include payment for product administration fees.

By signing above, the patient hereby certifies to the following:

I certify that I am not enrolled in any Federal health care program (such as Medicare, Medicaid, Veteran Affairs health care programs, Department of Defense health care programs, TRICARE, or CHAMPUS) If I am enrolled in the Commercial Bridge Program, I certify that all information provided herein is correct and complete, to the best of my knowledge. I acknowledge that any product provided no a complimentary basis. I certify that I will not submit or cause to be submitted any claims for payment or reimbursement for such products to any third-party payor, including any Federal health care programs. If I am or become in possession of such product, I understand that such product is only for me and I will not sell, trade, or distribute or otherwise transfer such product.

I understand that if I am enrolled in the Commercial I understand that if I am enrolled in the Commercial Bridge Program, any future changes to prescription drug coverage, or insurance coverage may affect whether I can continue to participate in the Commercial Bridge Program. I agree to contact KabiCare at 1-833-KABICARE (1-833-522-4227) and tell them about any changes to my prescription drug coverage, or insurance coverage, understand there is no purchase requirement associated with assistance through by or through the Commercial Bridge Program. I understand that completing this form does not guarantee that assistance will be provided.

Section 5: Fair Credit Reporting Act (FCRA) - Patient

The information I have provided is complete and accurate and will be used to decide if I am eligible to participate in the KabiCare Patient Assistance Program (the "PAP"). I agree that submitting my application is not a quarantee that I am entitled to participate in the PAP or that the PAP is obligated to provide me with any assistance. I understand that the PAP can revise, change, or terminate the program at any time. I authorize my healthcare providers and my health plan or insurers to give my medical and financial information to KabiCare, which administers the PAP on behalf of Fresenius Kabi, the distributor of the medicines, and to Experian and CoverMyMeds LLC, which assesses my income and ability to pa!, I authorize KabiCare and its service providers to obtain credit reports about me from one or more credit reporting agencies in order to verify my information and determine my eligibility to participate in the PAP. I authorize KabiCare and Experian to review The information I have provided is complete and accurate in the PAP. I authorize KabiCare and Experian to review my medical and financial information and to use it only to determine if I am eligible to participate in the PAP, to operate the PAP, or as otherwise required or permitted I law. Lunderstand and agree that KabiCare and Experian may contact me directly to verify the information I have submitted or to ask for additional information or documentation to process my application.

Section 6: Prescriber Certification and Statement of Medical Necessity

Prescriber Declaration: My signature certifies that the person named on this form is my patient, I will be supervising this patient's treatment, and the information that has been provided is complete and accurate to the best of my knowledge.

I also certify that I have made the clinical judgement that any products provided through kabiCare Patient Support Program(s), including KabiCare's Patient Assistance Program (the "PAP") are medically necessary and appropriate for the patient named on this form and will be used only by that patient. I will not use any such product or prescribe, provider, furnish, or dispense any portion thereof to any other person or patient. If I am or become in possession of such medications, I will not sell, resell, offer for sale, trade, or barter such products.

In addition, I certify that no claim for payment or reimbursement for any product furnished through KabiCare Patient Support Program(s), including the PAP, will be submitted to any third-party payer, including any Federal health care program (such as Medicare, Medicaid, Veteran Affairs health care programs, Department of Defense health care programs, TRICARE, or CHAMPUS), any other health care benefit plan, payer or patient, or returned for credit. In addition, I certify that no claim for payment or

I further certify that (a) any reimbursement investigation support or assistance provided to patients through KabiCare Patient Support Program is not made in exchange, directly or indirectly, for any past, present, or future recommendation, prescription, purchase, or use of the above therapy or any other product or service for or from anyone and (b) my decision to prescribe product was based solely on my determination of medical necessity as set forth herein. I understand that completing this form does not aurantee that assistance will be provided to does not guarantee that assistance will be provided to my patient.

Prescriber Acknowledgement: Where required by applicable law, regulation, or other applicable authority, I have obtained appropriate written authorization from the patient ("Legal Permission") permitting me to use and disclose my patients' health, demographic, and other individually identifiable information, including insurance and financial information, to Fresenius Kabi, its affiliates, its program administrator, and their respective agents, service providers and field reimbursement professionals for the purpose of respections the actions. service providers and field reimbursement professionals for the purpose of assessing the patients' insurance coverage and eligibility for participation in the KabiCare Patient Support Program(s), providing patient support programs, copay assistance, patient assistance, and/or reimbursement support in connection with the patient's treatment with product. I maintain records of such Legal Permission consistent with applicable law.

I appoint KabiCare on my behalf, to convey this prescription to the appropriate dispensing entity, to the extent permitted under applicable state law. Special Note: Prescribers in all states must follow applicable laws for a valid prescription. For prescribers in states with official prescription form requirements, please submit an actual prescription along with this enrollment form.

Section 7: Patient Assistance Program - Patient

Section 7: Patient Assistance Program - Patient Attestation

I certify that all information provided herein is correct and complete, to the best of my knowledge. To the extent I receive amy free product through programs offered by or through the KabiCare Patient Support Programs, including the KabiCare Patient Assistance Program ("PAP"), I acknowledge that such product is provided on a complimentary basis. I certify that I will not submit or cause to be submitted any claims for payment or reimbursement Products on any third-partip payor, including the Program of the Product of the Programs payor, in the Program of the Programs of the Product is only for me, and I will not give such product to anyone else. I agree that I will not sell; resell, offer for sale, trade, barter, distribute or otherwise transfer such product, I understand that if I am enrolled in a Medicare Part D Plan (including a Medicare Advantage Prescription Drug Plan) or other Federal health care program, I may not apply any assistance I receive to my "True Out of Pocket" ("TrODP") expenditures, and that it is my responsibility to notify such Federal health care program(s) of any assistance I may receive, including my enrollment in the PAP. I understand that if I am enrolled in the PAP, any future changes to my income, prescription drug coverage, or insurance coverage may affect whether I can continue to participate in the PAP. I agree to contact KabiCare at 1-833-KABICARE (1-833-522-4227) and inform them about any changes to my income, prescription drug coverage, or insurance coverage understand there is no them about any changes to my income, prescription drug overage, or insurance coverage. I understand there is no purchase requirement associated with assistance through by or through any KabiCare Patient Support Program(s), including the PAE1 understand that completing this form does not guarantee that assistance will be provided.

I understand that Fresenius Kabi reserves the right at any I understand that resembles and reserves the right at any time and without notice to me to modify and/or discontinue any or all support offered by or through KabiCare Patient Support Programs, including modification of eligibility criteria, covered medications and immediate termination of assistance provided through the PAP.

